



Windows to the Brain: Introduction to Circuits

**Cortical Association
Prefrontal-Subcortical
Papez
Brainstem**

Katherine Taber, PhD, FANPA

MIRECC Assistant Director - Education

Research Health Scientist

W.G. "Bill" Hefner VAMC, Salisbury NC

Research Professor, Div Biomedical Sciences

Edward Via College of Osteopathic Medicine

Robin Hurley, MD, FANPA

MIRECC Associate Director - Education

ACOS/Research and Academic Affairs Service Line

W.G. "Bill" Hefner VAMC, Salisbury NC

Professor, Psychiatry and Radiology

Wake Forest School of Medicine

Source: <http://www.mirecc.va.gov/visn6/Tools-Tips.asp>

Use of text, images and other content are subject to the following terms and conditions:

Fair Use Is Permitted

Fair use of copyrighted material includes the use for non-commercial educational purposes, such as teaching, scholarship, research, criticism, commentary, news reporting, and other content. Unless otherwise noted, users who wish to download or print text and image files from this Web site for such uses may do so without the VISN 6 MIRECC's express permission, provided that they comply with the following conditions:

The content may only be used for personal, educational or non-commercial purposes;


Users must always specifically cite the author(s) and source of the content every time the material is used, as they would for material from any printed work;

None of the content may be altered or modified.

Warranty

By downloading, printing, or otherwise using text and image files from this website, users agree and warrant that they will limit their use of such files to fair use.

Major Long Cortical Association Tracts



Brief guide to neuropsychiatric symptoms associated with injury to tract.*

superior fronto-occipital (subcallosal) fasciculus

orbital & medial prefrontal cortex ↔ parietal cortex

akinetic mutism; disordered initiation & preparation of speech movements; transcortical motor aphasia; anomia & reduction of spontaneous speech with normal articulation

cingulum - short fibers

cingulate cortex ↔

frontal, parietal, occipital & temporal cortex

cingulum - long fibers

frontal cortex ↔

temporal cortex

Anterior - lack of emotional affective response to pain; anxiety; OCD; depression; panic; akinetic mutism **Posterior** - impaired integration of visuospatial & memory processing

superior longitudinal (arcuate) fasciculus

frontal cortex ↔ parietal, occipital & temporal cortex

R - left hemispatial neglect; **L** - conduction aphasia (fluent aphasia with impaired repetition, mostly preserved language comprehension); ideational apraxia (can't carry out skilled movements and/or commands); depression; speech arrest; anomia; **Posterior** - transcortical sensory aphasia (impaired auditory comprehension, intact repetition & fluent speech)

uncinate fasciculus

orbital & polar prefrontal cortex ↔ anterior temporal cortex

deficits in retrieval of past information: **R** - episodic context-dependent memory, personal experiences, autobiographical; **L** - context-free memory, general knowledge of facts

inferior fronto-occipital fasciculus

ventrolateral & dorsolateral prefrontal cortex ↔ posterior temporal & occipital cortex

R>L - impaired orienting of attention; visual recognition abnormalities; **R+L** - impaired pursuit eye movements; inaccurate reaching under visual guidance; impaired motion perception; **R or R+L** - impaired seeing/selecting in crowds; impaired spatial relations; visual agnosia & poor visual memory; impaired recognition of places & directions to get there; getting lost

inferior longitudinal fasciculus

temporal pole ↔ occipital cortex

disorders in recognition (visual agnosia) impaired visual recent memory; **R or R+L** - impaired face recognition (prospagnosia), visual object agnosia, visual hypoemotionality if cue presented visually; **R+L or L>R** - contralateral deficit in color vision (hemiachromatopsia); **L-bilateral** misnaming of objects presented by touch (tactoverbal dysfunction)

*Taber KH, Hurley RA. J Neuropsychiatry Clin Neurosci 2007;19(2):100-104.


Major Prefrontal - Subcortical Circuits

In psychiatry, the prefrontal cortex is generally divided into three principal areas. Each area has reciprocal connections with subcortical structures that form cortico-subcortical circuits.*

frontal temporal parietal occipital limbic


Dorsolateral circuit

mediates executive functions such as organization, planning & attention



Orbitofrontal circuit
mediates socially appropriate behavior, impulse control & empathy

Anterior cingulate circuit
produces motivation by balancing the inhibitory input of the supplemental motor area with its own stimulus that supports wakefulness & arousal



*Bonelli RM, Cummings JL. Dialogues Clin Neurosci. 2007;9(2):141-51.

Major Prefrontal - Subcortical Circuits


In psychiatry, the prefrontal cortex is generally divided into three principal areas. Each area has reciprocal connections with subcortical structures that form cortico-subcortical circuits.

These prefrontal-subcortical circuits are formed by chains of neurons with cell bodies in gray matter structures (both cortical and subcortical) connected by the axons which form the white matter. Recently the evidence supporting a similar reciprocal circuit to the cerebellum has strengthened, although its functions are still controversial.

Dorsolateral circuit
mediates executive functions such as organization, planning and attention


Orbitofrontal circuit
mediates socially appropriate behavior, impulse control, and empathy

Anterior Cingulate circuit
produces motivation by balancing the inhibitory input of the supplemental motor area with its own stimulus that supports wakefulness and arousal




Circuit of Papez

A schematic diagram of the emotion and memory circuit of Papez is color-coded to match the summary of subcortical structures and the sectional atlases. The location and extent of the Anterior Cingulate cortex (light gold), fornix and mammillary body are indicated on a midline sagittal magnetic resonance image. The locations of the remaining structures are shown on a coronal magnetic resonance image.




Brainstem - Amine Neurotransmitters

These small brainstem nuclei project very widely in the brain. They provide essential modulation of the brain systems that subserve multiple functions including behavior, cognition and mood. The approximate locations and extents of nuclei that are important sources for a particular neurotransmitter are color-coded onto a sagittal magnetic resonance image.*




*Hurley RA, Flashman LA, Chow TW, Taber KH. J Neuropsychiatry Clin Neurosci 2010;22(1):iv, 1-6.


Neurotransmitters - Acetylcholine



Neurotransmitters - Serotonin



Neurotransmitters - Norepinephrine



Neurotransmitters - Dopamine

