

VHA OFFICE OF MENTAL HEALTH AND SUICIDE PREVENTION

Providing SST 1 on 1

June 26, 2023

Elizabeth Gilbert, Ph.D.

Rebecca Pasillas, Ph.D.

VA Social Skills Training for Serious Mental Illness (SST) Training Program
VISN 5 Mental Illness Research, Education and Clinical Center (MIRECC)



Choose **VA**

VA



U.S. Department
of Veterans Affairs

Acknowledgments

- We would like to thank Melanie Bennett, Ph.D. and Jennifer Aakre, Ph.D., for their contributions to previous versions of this presentation.



TMS

- If you would like to get continuing education credit (CE) for attending this presentation, you need to have registered in TMS prior to the presentation. After this presentation, you also need to confirm in TMS that you attended the event and to complete the TMS course survey within 15 days after the presentation.
- Even if you do not wish to receive CE credit, please complete the TMS course survey so that we may receive program evaluation feedback and continue to offer CE for our training events.
- We encourage you to complete the TMS course survey immediately after the presentation today.



Agenda

- Review previous research on social skills training implemented in-person, 1 on 1
- Discuss ideas about how best to implement the Bellack et al. SST model in in-person individual sessions



Getting to know our audience

- Who on the call has implemented SST in-person 1 on 1?
- We have discussion time at the end to hear how your experiences relate to the information we present.



Choose **VA**

VA



U.S. Department
of Veterans Affairs

Evidence Base

- There is limited research on social skills training approaches implemented with people with serious mental illness on a 1 on 1 basis. It is not currently considered an EBP. However, the research that has been completed provides support that this type of intervention is likely helpful for individuals diagnosed with schizophrenia.
- There are studies (Hogarty, et. al, 1986 and 1991; Pratt, et al, 2017) that examined social skills training delivered in an individual session. Results were positive; they reported lower relapse rates than control group (Hogarty studies) and improved psychosocial functioning (Pratt study).



Choose **VA**

VA



U.S. Department
of Veterans Affairs

Hogarty et al., 1986 and 1991

- These studies do not solely focus on SST delivered 1 on 1
- In these studies, individually delivered social skills training (not exactly the Bellack et al. model but similar teaching strategies) was delivered in combination with other interventions.
- These studies provide some indirect support for the idea that individualized social skills training can be beneficial for individuals diagnosed with schizophrenia and schizoaffective disorders.
- Analysis of the positive outcomes (including relapse rates) of the combined interventions suggest that individualized social skills training can be a valuable component of a comprehensive treatment plan for schizophrenia and schizoaffective disorders.



Choose **VA**

VA



U.S. Department
of Veterans Affairs

HOPES-I (Pratt et al. 2017)

- Helping Older People Experience Success – Individually Tailored (HOPES-I) intervention
- Results: “Participants with baseline impairments in overall functioning and in each of the skill areas targeted by the program demonstrated significant improvements on related outcome measures.”
- No control group in study
- Intervention similar in some ways to SST but in a 1:1 format:
 - At least 1 recovery-oriented goal is set
 - Focus is on skill-building and role play practice
 - Curriculum of skills selected relevant to needs of client
- Additional element is assisting the client with generalizing skills in the community, “direct facilitation and support of goals and objectives (e.g., going to a fitness center with a participant to encourage independent access of this resource)” (p. 382)



VA EBPs and Older Veterans

- Older Veterans are significantly under-represented as EBP training cases
- Many older adults prefer psychotherapy to medication (e.g., Luck-Sikorski et al., 2017; Mohlman, 2012)
- Social Skills Training (SST; Bellack et al., 2004) is an important intervention to consider for older adults (i.e., ages 50 and older) with serious mental illness (SMI) given the evidence that social skill informed interventions demonstrate improved social and overall functioning in older adults with SMI (Jeste and Maglione, 2013).
- Consider offering 1 on 1 SST to an older Veteran



IVAST

- In Vivo Amplified Skills Training (IVAST) (Glynn, et al., 2002; Liberman, et al., 2002)
- Clients participate both in a social skills training group (UCLA Social and independent Living Skills Program) and also have a case manager who helps them implement social skills in the community.
- Mean age of the Veterans in this study was approximately 43 years old.
- The case managers help the clients adapt the social skills to their environments.
- The case manager offers encouragement and reinforcement for successful generalization of skills to the community.



Choose **VA**

VA



U.S. Department
of Veterans Affairs

IVAST

- An evaluation of IVAST (Glynn, et al., 2002) reports that compared to participants attending SST groups alone, participants in IVAST condition over a 60 week period had:
 - higher levels of interpersonal problem-solving skills
 - greater social adjustment
 - better quality of life



SST & Individual Coaching

- Pilot study: Social skills training group (Bellack et al., 1997 model) with supplementary individual coaching in a rural community mental health center over 8-months (Gottlieb, et al., 2005)
- “Skills coaches” provided extra skill practice in community settings; weekly 30-40 minute sessions
- Compared social skills group with supplementary coaching to social skills group without additional coaching
- Assessments at baseline, 4 months, and 8 months



Choose **VA**

VA



U.S. Department
of Veterans Affairs

SST & Individual Coaching

- Group 1 (no skills coach)
 - Therapist ratings: no improvement in skills, attentiveness, & cooperation at 4- and 8-months
 - Social functioning: improvement at 4 months, decreased at 8 months
- Group 2 (skills coach)
 - Therapist ratings: improvement in skills, attentiveness, & cooperation
 - Social functioning: improvements at 4 months and continued at 8 months
 - Skills coaching ratings: skill performance improved at 4- and 8-months



Bellack et al. SST

- Individual social skills training “Can be used either to supplement group-based training or alone” (Bellack et al., p. 75)
- Single-case-study using multiple-baseline design demonstrate improvement in social skills (Bellack et al., pgs. 19-20)
- Bellack, A.S., Mueser, K.T., Gingerich, S., & Agresta, J. (2004). *Social Skills Training for Schizophrenia: A Step-by-Step Guide*. New York: Guilford Press.



Choose **VA**

VA



U.S. Department
of Veterans Affairs

Why Do Individual SST?

- Useful for
 - People who are reluctant to come to a group – to prepare them to join the group
 - People who have difficulty participating in a group setting (e.g., reluctant to be in a group setting after the pandemic)
 - People who may have specific cognitive, functional, and/or sensory limitations who may benefit from an individualized approach
 - People who can't make the SST group time
 - Tackling personalized situations in order to increase generalization of skills to the community
 - People who want to do intensive work on a specific social skills domain
 - Creating more opportunity to support the generalization of skills in the community situations in which they will be used
 - The maintenance of skills following SST group participation
 - Enhancing other types of psychotherapy (e.g., CBT, Interpersonal Therapy)



Format of Individual SST

- Format may depend on whether the individual SST is a supplement to the group or alone.
- Start with an introductory meeting
 - Review rationale of individual SST
 - Review goal(s) and format of sessions
- Duration and frequency of the individual SST sessions depends on the individual's goals.
- One approach is to start with 3-5 meetings and then reevaluate based on progress the frequency and duration of planned sessions. Individuals can return for boosters or 5-session blocks as situations in their lives evolve.



What's Done in Individual SST

- Content depends on purpose
- Can focus on gearing up to integrate with an existing SST group
 - If group is ongoing, make sure the client in the individual sessions receives training on the same skills that are taking place in the group so they will be up to speed when rejoining.
- Can focus on specialized skills based on an individual's needs and preferences
- Can focus on modifying the steps of the skill so that it fits the social, cultural, and linguistic context of the situation.
- Can work through a hierarchy of social skills and role plays, gradually progressing from simple skills to more complex skills and role plays
- Can consider doing more generalization work (i.e., coaching clients to practice their skills in real-life settings in their community)
- May be more practical to involve the family and others in the client's environment when SST is done 1 on 1. May want to educate the family and encourage family support and participation in supporting the client's use of SST skills with the consent of the client.



Choose  VA

VA



U.S. Department
of Veterans Affairs

Use the SST Session Sequence for Individual Sessions

Welcome and set agenda

1. Review outside practice assignments (for sessions other than the first session)
2. Establish a rationale for the skill
3. Briefly have ~~members~~ **the client** share a relevant experience or rationale
4. Explain the steps of the skill
5. Model the skill; review with ~~members~~ **the client**
6. Have a ~~group member~~ **the client** role play
7. Elicit and provide positive, then constructive feedback
8. Have the ~~member~~ **the client** role play again
9. Elicit and provide positive, then constructive feedback
10. Repeat role play again and elicit/provide positive feedback
11. Repeat Steps 6-10 with ~~each group member~~ **the client**. ***The individual client can do more than one set of role plays in a 50 minute session.**
12. Develop outside practice assignments



Choose **VA**

VA



U.S. Department
of Veterans Affairs

Examples of Individual SST that is Personalized to a Client's Needs and Goals

- Individual wants to be more assertive with their live-in significant other
 - Work on refusing requests, expressing angry feelings, compromise and negotiation
- Individual has symptoms of depression and feelings of worthlessness that kept them from participating in SST groups
 - Work on skills for starting and maintaining conversations, connecting social skills with improved mood
- Young adult, wants to get a job
 - Work on vocational/work skills



Individual SST on a longer-term basis

- Suggested enhancements
 - To avoid repetition, invite others to serve as the role play buddy in role plays
 - Practice skills in the community
 - Add content as needed to put SST in the context of the client's life experience. Can include psychoeducation, stress management, methods for coping with persisting sx., etc. and how SST helps with all of these.
- SST can either be the main intervention or one of the strategies a provider employs in individual work.



Choose **VA**

VA



U.S. Department
of Veterans Affairs

SST in Individual Psychotherapy

- “While cognitive behavior therapy also has become an evidence-based treatment for residual psychotic symptoms, few clinicians or investigators have recognized the importance of social skills in this modality's reliance on behavioral assignments and “experiments.” Social skills training is implicitly involved in instigating favorable outcomes in both cognitive behavior therapy and interpersonal therapy, but more explicit integration and adaptation of skills training methods in these therapies holds promise for improving their impact in schizophrenia”
- From Kopelowicz, A., Liberman, R.P., & Zarate, R. (2006). Recent Advances in Social Skills Training for Schizophrenia. *Schizophrenia Bulletin*, 32(S1), S12–S23, p S21.



Choose  VA

VA



U.S. Department
of Veterans Affairs

Q & A and Discussion



Choose **VA**

VA



U.S. Department
of Veterans Affairs

Reach out

- Elizabeth Gilbert, Ph.D., Program Manager VA SST Training Program elizabeth.gilbert@va.gov
- Rebecca Pasillas, Ph.D., Program Evaluator/National Trainer and Consultant, VA SST Training Program rebeccamelissa.pasillas@va.gov



References

- Jeste, D. V., & Maglione, J. E. (2013). Treating older adults with schizophrenia: Challenges and opportunities, *Schizophrenia Bulletin*, 39(5), 966–968. <https://doi.org/10.1093/schbul/sbt043>
- Glynn, S. M., Marder, S. R., Liberman, R. P., Blair, K., Wirshing, W. C., Wirshing, D. A., Ross, D., & Mintz, J. (2002). Supplementing clinic-based skills training for schizophrenia with manual-based community support: Effects on social adjustment of patients with schizophrenia. *American Journal of Psychiatry*, 159(5), 829–837. <https://doi.org/10.1176/appi.ajp.159.5.829>
- Gottlieb, J. D., Pryzgoda, J., Neal, A., & Schulberg, D. (2005). Generalization of skills through the addition of individualized coaching: Development and evaluation of a social skills training program in a rural setting. *Cognitive and Behavioral Practice*, 12(3), 324–338. [https://doi.org/10.1016/S1077-7229\(05\)80055-0](https://doi.org/10.1016/S1077-7229(05)80055-0)
- Hogarty, G. E., Anderson, C. M., Reiss, D. J., Kornblith, S. J., Greenwald, D. P., Javna, C. D., & Madonia, M. J. (1986). Family psychoeducation, social skills training, and maintenance chemotherapy in the aftercare treatment of schizophrenia. I. One-year effects of a controlled study on relapse and expressed emotion. *Archives of General Psychiatry*, 43(7), 633-642. <https://doi.org/10.1001/archpsyc.1986.01800070019003>
- Hogarty, G. E., Anderson, C. M., Reiss, D. J., Kornblith, S. J., Greenwald, D. P., Javna, C. D., & Madonia, M. J. (1991). Family psychoeducation, social skills training, and maintenance chemotherapy in the aftercare treatment of schizophrenia. II. Two-year effects of a controlled study on relapse and adjustment. *Archives of General Psychiatry*, 48, 340-347. <https://doi.org/10.1001/archpsyc.1991.01810280056008>
- Liberman, R. P., Glynn, S., Blair, K. E., & Marder, S. R. (2002). In vivo amplified skills training: Promoting generalization of independent living skills for clients with schizophrenia. *Psychiatry*, 65(2), 137–155. <https://doi.org/10.1521/psyc.65.2.137.19931>
- Luck-Sikorski, C., Stein, J., Heilmann, K., Maier, W., Kaduszkiewicz, H., Scherer, M., Weyerer, S., Werle, J., Moor, L., Bock, J. O., König, H.H., & Riedel-Heller, S. (2016). Treatment preferences for depression in the elderly. *International Psychogeriatrics*, 29(3), 389-398. <https://doi.org/10.1017/S1041610216001885>
- Mohlman J. (2012). A community based survey of older adults' preferences for treatment of anxiety. *Psychology and Aging*, 27(4), 1182–1190. <https://doi.org/10.1037/a0023126>
- Pratt, S.I., Mueser, K.T., Wolfe, R., Santos, M. M., & Bartels, S. J. (2017). One size doesn't fit all: A trial of individually tailored skills training. *Psychiatric Rehabilitation Journal*, 40(4), 380-386. <http://doi.org/10.1037/prj0000261>



TMS

- If you would like to get continuing education credit (CE) for attending this presentation, you need to have registered in TMS prior to the presentation. After this presentation, you also need to confirm in TMS that you attended the event and to complete the TMS course survey within 15 days after the presentation.
- Even if you do not wish to receive CE credit, please complete the TMS course survey so that we may receive program evaluation feedback and continue to offer CE for our training events.
- We encourage you to complete the TMS course survey immediately after the presentation today.

